

Resumption of Pulmonary Function Testing during the Post-Peak Phase of the COVID-19 Pandemic

A Position Statement from the Canadian Thoracic Society and the Canadian Society of Respiratory Therapists

Sanja Stanojevic^a, François Beaucage^b, Vikram Comondore^c, Marie Faughnan^d, Tom Kovesi^e, Carolyn McCoy^f, Colm McParland^{g,k}, David Pawluski^h, Farzad Refahiⁱ, Jeremy Road^j, Micah Kooperberg^k

^aDepartment of Community Health and Epidemiology, Dalhousie University, Halifax, NS, Canada; ^bHôpital du Sacré-Coeur de Montréal and l'Université de Montréal, Montréal, QC, Canada; ^cRespiratory and Sleep Medicine, William Osler Health System, Department of Medicine, McMaster University, Hamilton, ON, Canada; ^dSt Michael's Hospital Unity Health Toronto, Division of Respiriology, Department of Medicine, University of Toronto, Toronto, ON, Canada; ^eDepartment of Pediatrics, Children's Hospital of Eastern Ontario, University of Ottawa, Ottawa, ON, Canada; ^fCanadian Society of Respiratory Therapists, Ottawa, Ontario, Canada; ^gDivision of Respiriology, Department of Medicine, Dalhousie University, Halifax, NS, Canada; ^hThe Canadian Association of Cardio-Pulmonary Technologists, University of Alberta Hospital, Edmonton, AB, Canada; ⁱMarkham-Stouffville Hospital, Markham, ON, Canada; ^jDivision of Respiratory Medicine, University of British Columbia, Vancouver, BC, Canada; ^kNova Scotia Health Authority, Halifax, NS, Canada.

Version 2.0 - December 9, 2020

This position statement aims to provide rapid guidance for resumption of pulmonary function services during the post-peak phase¹ of the COVID-19 pandemic (i.e., period of low community prevalence). If the prevalence of COVID-19 increases (pandemic phase – high community prevalence) or decreases (post-pandemic phase – controlled) these recommendations may need to be revised. **These recommendations are based on the consensus of the authors, who are members of either the Canadian Thoracic Society or the Canadian Society of Respiratory Therapists and are subject to change as information regarding COVID-19 and its effects are further understood.** We plan to update this guidance as new information becomes available and recommend periodically checking the Canadian Thoracic Society website (<https://cts-sct.ca/covid-19/>) or Canadian Society of Respiratory Therapists website (<https://www.csrt.com/position-statements/>) for updates. The Canadian Association of Cardio-Pulmonary Technologists endorses this position statement.

Pulmonary function tests (PFTs) are important for the diagnosis, management and monitoring of disease and there is an urgent need to safely resume services. Patients cannot wear a mask during the test and physical distancing of two meters between the patient and the respiratory therapist/pulmonary function technologist performing PFTs is not always feasible. Patients must also come in contact with laboratory equipment. Coughing induced during and after a PFT may result in high aerosol generation², which can spread droplets from an infected individual even if they are asymptomatic. Furthermore, many patients who attend pulmonary function laboratories have underlying respiratory conditions that render them at higher risk of complications from COVID-19.³ Given these factors, the safety of both patients and staff require careful consideration when resuming PFT services.

We recognize the diversity of circumstances in different jurisdictions across Canada, and between health facilities. Guidance from the Public Health Agency of Canada and from local public health or infection

control units regarding health care facility capacity to resume services, public health measures and screening practices should supersede this document. We recommend that facility specific resumption plans include consultation with a multi-disciplinary team of relevant stakeholders. This could include a respiratory therapist/pulmonary function technologist, respirologist, medical director, manager, infection control representative, administrative team member including registration and scheduling personnel, screening team, environmental services, facilities, biomedical engineering, filter/equipment providers and/or occupational health, as appropriate to the facility.

Current evidence suggests that SARS-CoV-2 is transmitted through droplets and aerosols⁴⁻⁸, typically when symptomatic individuals cough or sneeze.⁸ There is also evidence that SARS-CoV-2 can be transmitted by asymptomatic individuals, including by speaking and coughing.⁶⁻¹³ PFT test manoeuvres can generate expiratory airflow rates that are higher than those during cough, and are high enough to aerosolise microorganisms.¹⁴ The risk of aerosol generation during PFTs will depend on both the test and patient. Methacholine challenge testing, exercise challenge testing, and cardiopulmonary exercise testing are higher risk than spirometry, six-minute walk test, and measurement of lung volumes or diffusion capacity. ***Given these factors, it is the consensus of the authors that an abundance of caution be used to protect patients and staff until there is clear evidence regarding the risk of transmission of SARS-CoV-2 during PFTs.*** This is consistent with guidance from other national and international organizations.¹⁵⁻¹⁷

Screening

- Screening of patients and PFT laboratory staff for COVID-19 is recommended.
 - Hospital based laboratories should follow the screening practices recommended by their facility and avoid duplicate and/or contradictory screening questionnaires
 - Independent laboratories can use screening forms developed by local public health offices.
- Screening should be performed for all patients scheduled to attend PFTs, within 72 hours of the scheduled test and upon entrance into the clinic or hospital on the day of testing. ***Patients who screen positive should not undergo PFTs.***
- Patients with pending COVID-19 test results should not have PFTs until COVID-19 can be ruled out.
- Patients who have tested positive for COVID-19 should NOT be permitted in the PFT laboratory until symptoms have resolved, and/or have two consecutive COVID-19 PCR swabs collected ≥ 24 hours apart that are negative.¹⁸ Viral shedding can occur after 10 days, therefore resolution of symptoms and a minimum period of 3 weeks after symptom onset is recommended when negative tests cannot be obtained.^{19, 20}
- Many patients who require PFTs also have chronic respiratory conditions with symptoms that are similar to those of COVID-19, therefore those with ***acute*** symptoms (e.g., self-reported and/or documented fever, cough, sore throat, dyspnea, or additional respiratory symptoms or myalgia or fatigue) should not be tested. In the absence of acute symptoms, or a change in symptoms, PFTs may be performed despite patients having respiratory symptoms.
- Staff with COVID-19 symptoms should not report to work and follow local occupational health policies regarding testing for COVID-19.
- Since COVID-19 screening and testing can lead to false negatives, it is prudent to assume that all individuals may be asymptomatic transmitters of COVID-19. This risk of false negatives should be considered in context of the level of community spread in each jurisdiction.

Personal Protective Equipment

- Based on the available evidence and the consensus opinion of our expert authors and other major professional bodies,^{15, 16, 21} during pandemic and post-peak pandemic phases, ***fit-tested N95 masks (or equivalent), face shields and protection appropriate for aerosol generating medical procedures (AGMP) should be employed by clinicians conducting PFTs.***
- We acknowledge that some PFTs are more likely to induce cough than others. Nonetheless, out of an abundance of caution, we recommend AGMP precautions be taken for all PFTs.

Testing Environment

- Negative pressure rooms or HEPA filtration systems with UV germicidal lamps are recommended while there is documented community spread of COVID-19 (pandemic phase/post peak phase).²² The use of these devices does not replace the need for appropriate PPE.
- When the use of negative pressure rooms, HEPA filtration and UV light are not available in the testing environment, pulmonary function testing may still resume, provided PPE, time-between patients and cleaning precautions are followed to protect patients and staff from viral transmission.
- Patients should be asked to wear procedure masks while in waiting areas and adequate physical distancing (2-meter distance) should be followed in waiting rooms. Staggered appointment times may be appropriate, to limit the number of patients in waiting areas. Where possible, one-way patient flow in laboratories should be implemented.
- Patients and staff should follow appropriate hand hygiene measures.
- Staff should be educated and trained on correct donning and doffing technique for PPE.
- Single use bacterial/viral inline filters that meet international standards of filtration performance should be used for each patient²³ (e.g. filters with minimum proven efficiency for high expiratory flow of 600 to 700 L/min). Laboratories should confirm with manufacturers that filters adequately protect against COVID-19 sized viral particles.
- Patients should be tested one at a time in designated enclosed testing rooms. Testing room door should remain closed for the duration of test.
- There should be sufficient time between appointments to allow for aerosols to settle or dissipate and for cleaning and disinfecting of surfaces. Time between patients should consider the number of air changes in the testing room (determined by each facility) to ensure 99.0 % - 99.9 % removal of airborne microorganisms. See CDC guidance for time required per air change rate.²⁴ For example, at 6 air exchanges per hour a minimum of 46 and 69 minutes is required for 99.0% and 99.9% removal, respectively. [Additional guidance that considers ambient relative humidity is available.](#)²⁵
- Facilities are encouraged to determine their air exchange rate in each PFT testing room. Where it is not feasible to determine the air exchange rate, a minimum of three hours should elapse between test termination and cleaning /subsequent testing.^{26, 27}
- Equipment used for testing and testing rooms should be easy to clean and disinfect between patients. The person performing the cleaning should wear the PPE recommended above. Cleaning and disinfecting should include wiping down all surfaces that the patient and staff came into contact with and that were within a 2-meter radius of the patient and staff member. A hospital-grade antiviral disinfectant should be used. All non-essential accessories should be removed from the designated testing room.

Scheduling and Patient Prioritisation

- Consistent with existing recommendations during the pandemic phase of COVID-19 only **essential** testing should be performed (i.e. where testing is required for immediate management decisions).^{15,28,29}
- During the post-peak phase, services should resume for patients in whom care decisions require PFTs (e.g. lung transplant or lung resection candidates, monitoring post-lung transplant, chemotherapy surveillance or pre/post monitoring of cancer therapy protocols associated with lung radiation damage, pre-operative risk evaluation, urgent clinical decision for patients with acute or chronic lung disease).
- Routine testing (i.e. tests that are not required to guide current clinical decisions) should be postponed until capacity increases to meet the demands of essential cases.
- During the pandemic and post-peak phases of the pandemic, PFT requests should be reviewed by a trained clinician (medical director/respirologist/lead respiratory therapist or pulmonary function technologist) to ensure that patients in whom PFT results are required to make care decisions are prioritized.
- Since modifications to the testing environment (e.g. requirement for PPE, extra cleaning etc.) are necessary, capacity for testing will be decreased. Time required to complete testing, including donning/doffing of PPE, cleaning/disinfecting and wait time between patients, will be considerably longer than what was previously possible. Scheduling of visits should consider these additional time demands.
- The logistics within each facility regarding the direction/spacing of patient flow should be considered to minimize contact between patients (e.g. waiting room capacity with adequate spacing). It may be beneficial to instruct patients to arrive no more than 10 minutes prior to their appointment time, to limit contact with staff and other patients.
- In selected patients with chronic respiratory conditions, spirometry and 6-minute walk tests may be performed by the patient at home where available (with appropriate guidance, training and quality control procedures), with results reviewed by the clinical team via telemedicine.

Additional Considerations

- Children with COVID-19 are more likely to be asymptomatic. While a child-friendly testing environment should be maintained as much as is feasible, at no time should appropriate infection control be overlooked to accomplish this.
- When required, one caregiver (e.g. parent; translator; or physical/mental/social support person) should be permitted to accompany the patient to the testing area. This caregiver must also be screened, follow hand hygiene procedures and be asked to wear a facemask.
- Patients and caregivers should be asked to wear masks while in the healthcare facility.

Resumption of Normal Services

- These recommendations should be maintained until public health determines that COVID-19 is controlled and in the post-pandemic phase.

References

1. World Health Organization. Pandemic Influenza Preparedness and Response: A WHO Guidance Document. Geneva: World Health Organization. 2009. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK143061/>. Accessed June 18, 2020.
2. [Greening NJ, Larsson P, Ljunström E, et al. Small droplet emission in exhaled breath during different breathing manoeuvres: Implications for clinical lung function testing during COVID-19. *Allergy*. 2020. <https://doi.org/10.1111/all.14596>.](#)
3. Jain V, Yuan J-M. Systematic review and meta-analysis of predictive symptoms and comorbidities for severe COVID-19 infection. *medRxiv* (2020). doi: 10.1101/2020.03.15.20035360.
4. [Prather KA, Wang CC, Schooley RT. Reducing transmission of SARS-CoV-2. *Science*. 2020. \[doi:10.1126/science.abc6197\]\(https://doi.org/10.1126/science.abc6197\).](#)
5. [Centers for Disease Control and Prevention. Scientific brief: SARS-CoV-2 and potential airborne transmission. 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/more/scientific-brief-sars-cov-2.html>. Accessed on December 9, 2020.](#)
6. [World Health Organization. Coronavirus disease \(COVID-19\): How is it transmitted? Posted on October 20, 2020. Available from: <https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-how-is-it-transmitted>. Accessed on December 9, 2020.](#)
7. [Government of Canada. COVID-19: Main modes of transmission. Posted on November 5, 2020. Available from: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/health-professionals/main-modes-transmission.html>. Accessed on December 9, 2020.](#)
8. [Santarpia JL, Rivera DN, Herrera VL, et al. Aerosol and surface contamination of SARS-CoV-2 observed in quarantine and isolation care. *Sci Rep* 10, 12732 \(2020\). <https://doi.org/10.1038/s41598-020-69286-3>.](#)
9. World Health Organization. Modes of transmission of virus causing COVID-19: implications for IPC precaution recommendations. Posted update on July 9, 2020. Available from: <https://www.who.int/news-room/commentaries/detail/modes-of-transmission-of-virus-causing-covid-19-implications-for-ipc-precaution-recommendations>. Accessed July 9, 2020.
10. Anderson EL, Turnham P, Griffin JR, et al. Consideration of the Aerosol Transmission for COVID-19 and Public Health. *Risk Anal*. 2020;40(5):902-907. doi:10.1111/risa.13500.
11. Asadi S, Bouvier N, Wexler AS, et al. The coronavirus pandemic and aerosols: Does COVID-19 transmit via expiratory particles? *Aerosol Sci Tech*. 2020;54(6):635-638. doi:10.1080/02786826.2020.1749229.
12. Dhand R, Li J. Coughs and Sneezes: Their Role in Transmission of Respiratory Viral Infections, Including SARS-CoV-2. *Am J Resp Crit Care Med*. 2020. doi:10.1164/rccm.202004-1263PP.
13. Long Q, Tang X, Shi Q, et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med*. 2020. doi:10.1038/s41591-020-0965-6.
14. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J*. 2005;26:319–338. doi: 10.1183/09031936.05.00034805.
15. European Respiratory Society. Recommendation from ERS Group 9.1 (Respiratory function technologists/Scientists) Lung function testing during COVID-19 pandemic and beyond. Available from: <https://ers.app.box.com/s/zs1uu88wy51monr0ewd990itoz4tsn2h>. Accessed June 18, 2020.
16. Association for Respiratory Technology and Physiology. ARTP COVID19 Update. Posted on March 18, 2020. Available from: <https://www.artp.org.uk/News/artp-covid19-update-18th-march-2020>. Accessed June 18, 2020.
17. Thomas JP, Srinivasan A, Wickramarachchi CS. Evaluating the national PPE guidance for NHS healthcare workers during the COVID-19 pandemic. *Clin Med*. 2020;20(3):242-247. doi: 10.7861/clinmed.2020-0143.
18. Public Health Agency of Canada. Infection prevention and control for COVID-19: Second interim guidance for acute healthcare settings. Posted on April 30, 2020. Available from: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/health-professionals/infection-prevention-control-covid-19-second-interim-guidance.html>. Accessed on June 12, 2020.
19. World Health Organization. Criteria for releasing COVID-19 patients from isolation. Posted on June 17, 2020. Available from: <https://www.who.int/news-room/commentaries/detail/criteria-for-releasing-covid-19-patients-from-isolation>. Accessed on June 18, 2020.

20. Sethuraman N, Jeremiah SS, Ryo A. Interpreting Diagnostic Tests for SARS-CoV-2. *JAMA*. 2020;323(22):2249–2251. doi:10.1001/jama.2020.8259.
21. National Institute for Occupational Safety and Health, Occupational Safety and Health Administration. Hospital Respirator Protection Program Toolkit. 2015. Available from: <https://www.osha.gov/Publications/OSHA3767.pdf>. Accessed June 18, 2020.
22. Ereth MH, Hess DH, Driscoll A. Particle control reduces fine and ultrafine particles greater than HEPA filtration in live operating rooms and kills biologic warfare surrogate. *Am J Infect Control*. 2020. doi.org/10.1016/j.ajic.2019.11.017.
23. Unstead M, Stearn MD, Cramer D, et al. An audit into the efficacy of single use bacterial/viral filters for the prevention of equipment contamination during lung function assessment. *Respir Med*. 2006;100(5):946-50. doi: 10.1016/j.rmed.2005.09.015.
24. Centers for Disease Control and Prevention. Guidelines for Environmental Infection Control in Health-Care Facilities: Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). 2020:223. Available from: <https://www.cdc.gov/infectioncontrol/guidelines/environmental/appendix/air.html>. Accessed June 18, 2020.
25. Hurlburt AW, DeKleer RJ, Bryce EA. Clearing the air: Addressing air clearance times for infectious aerosols in healthcare facilities. *Infect Control Hosp Epidemiol*. 2020;1-2. doi: 10.1017/ice.2020.432.
26. VanDoremalen N, Bushmaker T, Morris DH, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*. 2020;382:1564-1567. doi: 10.1056/NEJMc2004973.
27. Bourouiba, L. Turbulent Gas Clouds and Respiratory Pathogen Emissions: Potential Implications for Reducing Transmission of COVID-19. *JAMA*. 2020;323(18):1837-1838. doi:10.1001/jama.2020.4756.
28. American Thoracic Society. Pulmonary Function Laboratories: Advice Regarding COVID-19. 2020. Available from: <https://www.thoracic.org/professionals/clinical-resources/disease-related-resources/pulmonary-function-laboratories.php>. Accessed June 18, 2020.
29. Canadian Society of Respiratory Therapists. Position Statement on Procedures Creating a Heightened Risk of Infection During an Outbreak of a Communicable Respiratory Disease. 2020. Available from: <https://www.csrt.com/wp-content/uploads/CSRT-Procedures-Duringan-Outbreak-April-2020-v2.pdf>. Accessed June 18, 2020.